


// DOSING AND ADMINISTRATION GUIDE

RYANODEX[®]

(dantrolene sodium)
for injectable suspension

FOR THE TREATMENT OF
MALIGNANT HYPERTHERMIA (MH)



INDICATIONS

RYANODEX[®] is indicated for the treatment of malignant hyperthermia in conjunction with appropriate supportive measures, and for the prevention of malignant hyperthermia in patients at high risk.

Please see Important Safety Information on back and enclosed full Prescribing Information.

 **RYANODEX[®]**
(dantrolene sodium)
for injectable suspension
250 mg per vial

EARLY RECOGNITION AND RAPID TREATMENT REDUCE MORBIDITY AND MORTALITY IN AN MH CRISIS¹

EVERY MINUTE COUNTS IN THE TREATMENT OF MH¹

For each 30-minute delay in the administration of dantrolene, the risk of MH-related complications increases 1.6 times.²

KNOW THE CLINICAL SIGNS OF MH

Malignant hyperthermia can happen at any time during anesthesia, or shortly after the discontinuation of a triggering agent.³ Known triggering agents include the volatile anesthetics (isoflurane, desflurane, sevoflurane) and succinylcholine.⁴ Signs and symptoms include, but are not limited to²:

- ▲ Masseter spasm
- ▲ Generalized muscle rigidity
- ▲ Hypercarbia
- ▲ Tachypnea
- ▲ Tachycardia
- ▲ Rapidly increasing temperature
- ▲ Hyperkalemia



**IF YOU SUSPECT MH, CONTACT THE 24-HOUR
MHAUS* EMERGENCY HOTLINE AT 800.644.9737**

*Malignant Hyperthermia Association of the United States.



**/ ONLY THE RYANODEX[®] FORMULATION OF
DANTROLENE SODIUM CAN BE ADMINISTERED
IN LESS THAN 1 MINUTE^{5,6} /**

IMPORTANT SAFETY INFORMATION

RYANODEX[®] is not a substitute for appropriate supportive measures in the treatment of malignant hyperthermia (MH), including discontinuing use of MH-triggering anesthetic agents, managing the metabolic acidosis, instituting cooling when necessary, and administering diuretics to prevent late kidney injury due to myoglobinuria (the amount of mannitol in RYANODEX[®] is insufficient to maintain diuresis).

**Please see Important Safety Information on back
and enclosed full Prescribing Information.**

RYANODEX®: 4 SIMPLE STEPS TO ADMINISTRATION IN LESS THAN 1 MINUTE^{5,6}



1. RECONSTITUTE

- Add 5 mL of sterile water for injection (without bacteriostatic agent)



2. SHAKE

- Shake vial to ensure an orange-colored uniform suspension*
 - Should take approximately 10 seconds⁷



3. FILL SYRINGE

- Visually inspect the vial for particulate matter and discoloration prior to administration
- Draw the appropriate dose of the reconstituted suspension into the syringe



4. ADMINISTER

- RYANODEX® (dantrolene sodium) for injectable suspension should be administered by intravenous push

*Must use the contents of the vial within 6 hours after reconstitution. Store reconstituted suspensions at controlled room temperature (68°F to 77°F or 20°C to 25°C).

RYANODEX®: REQUIRES FEWER VIALS RECONSTITUTED AND SIGNIFICANTLY LESS VOLUME ADMINISTERED THAN OTHER FORMULATIONS⁶⁻⁹

DOSE AND VOLUME NEEDED BASED ON PATIENT'S WEIGHT AND MHAUS RECOMMENDED DOSE OF 2.5 MG PER KG^{*10}

Patient's weight in kg	Patient's weight in pounds	MHAUS recommended dose (2.5 mg/kg)	Number of 250-mg RYANODEX® vials to reconstitute	Volume of reconstituted RYANODEX® to administer
5 kg	11 lbs	12.5 mg	1 vial	0.25 mL
10 kg	22 lbs	25 mg	1 vial	0.5 mL
15 kg	33 lbs	37.5 mg	1 vial	0.75 mL
20 kg	44 lbs	50 mg	1 vial	1 mL
25 kg	55 lbs	62.5 mg	1 vial	1.25 mL
30 kg	66 lbs	75 mg	1 vial	1.5 mL
35 kg	77 lbs	87.5 mg	1 vial	1.75 mL
40 kg	88 lbs	100 mg	1 vial	2 mL
45 kg	99 lbs	112.5 mg	1 vial	2.25 mL
50 kg	110 lbs	125 mg	1 vial	2.5 mL
55 kg	121 lbs	137.5 mg	1 vial	2.75 mL
60 kg	132 lbs	150 mg	1 vial	3 mL
65 kg	143 lbs	162.5 mg	1 vial	3.25 mL
70 kg	154 lbs	175 mg	1 vial	3.5 mL
75 kg	165 lbs	187.5 mg	1 vial	3.75 mL
80 kg	176 lbs	200 mg	1 vial	4 mL
85 kg	187 lbs	212.5 mg	1 vial	4.25 mL
90 kg	198 lbs	225 mg	1 vial	4.5 mL
95 kg	209 lbs	237.5 mg	1 vial	4.75 mL
100 kg	220 lbs	250 mg	1 vial	5 mL
105 kg	231 lbs	262.5 mg	2 vials	5.25 mL
110 kg	243 lbs	275 mg	2 vials	5.5 mL
115 kg	254 lbs	287.5 mg	2 vials	5.75 mL
120 kg	265 lbs	300 mg	2 vials	6 mL
125 kg	276 lbs	312.5 mg	2 vials	6.25 mL
130 kg	287 lbs	325 mg	2 vials	6.5 mL
135 kg	298 lbs	337.5 mg	2 vials	6.75 mL
140 kg	309 lbs	350 mg	2 vials	7 mL
145 kg	320 lbs	362.5 mg	2 vials	7.25 mL
150 kg	331 lbs	375 mg	2 vials	7.5 mL

Please see Important Safety Information on back and enclosed full Prescribing Information.

Dosing table continued

Patient's weight in kg	Patient's weight in pounds	MHAUS recommended dose (2.5 mg/kg)	Number of 250-mg RYANODEX® vials to reconstitute	Volume of reconstituted RYANODEX® to administer
155 kg	342 lbs	387.5 mg	2 vials	7.75 mL
160 kg	353 lbs	400 mg	2 vials	8 mL
165 kg	364 lbs	412.5 mg	2 vials	8.25 mL
170 kg	375 lbs	425 mg	2 vials	8.5 mL
175 kg	386 lbs	437.5 mg	2 vials	8.75 mL
180 kg	397 lbs	450 mg	2 vials	9 mL
185 kg	408 lbs	462.5 mg	2 vials	9.25 mL
190 kg	419 lbs	475 mg	2 vials	9.5 mL
195 kg	430 lbs	487.5 mg	2 vials	9.75 mL
200 kg	441 lbs	500 mg	2 vials	10 mL
205 kg	452 lbs	512.5 mg	3 vials	10.25 mL
210 kg	463 lbs	525 mg	3 vials	10.5 mL
215 kg	474 lbs	537.5 mg	3 vials	10.75 mL
220 kg	485 lbs	550 mg	3 vials	11 mL
225 kg	496 lbs	562.5 mg	3 vials	11.25 mL
230 kg	507 lbs	575 mg	3 vials	11.5 mL
235 kg	518 lbs	587.5 mg	3 vials	11.75 mL
240 kg	529 lbs	600 mg	3 vials	12 mL
245 kg	540 lbs	612.5 mg	3 vials	12.25 mL
250 kg	551 lbs	625 mg	3 vials	12.5 mL
255 kg	562 lbs	637.5 mg	3 vials	12.75 mL
260 kg	573 lbs	650 mg	3 vials	13 mL
265 kg	584 lbs	662.5 mg	3 vials	13.25 mL
270 kg	595 lbs	675 mg	3 vials	13.5 mL
275 kg	606 lbs	687.5 mg	3 vials	13.75 mL
280 kg	617 lbs	700 mg	3 vials	14 mL
285 kg	628 lbs	712.5 mg	3 vials	14.25 mL
290 kg	639 lbs	725 mg	3 vials	14.5 mL
295 kg	650 lbs	737.5 mg	3 vials	14.75 mL
300 kg	661 lbs	750 mg	3 vials	15 mL

- ▲ To calculate the volume in mL, multiply the patient weight in kg by desired dose (minimum dose 1 mg/kg; MHAUS recommended dose 2.5 mg/kg) and divide the product (mg dose) by 50.

*Calculated using 250-mg vials of RYANODEX® reconstituted with 5 mL of sterile water for injection USP (without a bacteriostatic agent). Labeled dose minimum of 1 mg/kg with a maximum cumulative dosage of 10 mg/kg. If the physiologic and metabolic abnormalities of MH reappear, repeat RYANODEX® dosing by intravenous push starting with 1 mg/kg.

/ CONTACT AN EAGLE REPRESENTATIVE TODAY AT 1-855-318-2170 TO ORDER RYANODEX® FOR YOUR FACILITY /

NDC 42367-540-32

To report SUSPECTED ADVERSE REACTIONS, contact Eagle Pharmaceuticals, Inc. at 1-855-318-2170 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

INDICATIONS

RYANODEX® is indicated for the treatment of malignant hyperthermia in conjunction with appropriate supportive measures, and for the prevention of malignant hyperthermia in patients at high risk.

IMPORTANT SAFETY INFORMATION

RYANODEX® is not a substitute for appropriate supportive measures in the treatment of malignant hyperthermia (MH), including discontinuing use of MH-triggering anesthetic agents, managing the metabolic acidosis, instituting cooling when necessary, and administering diuretics to prevent late kidney injury due to myoglobinuria (the amount of mannitol in RYANODEX® is insufficient to maintain diuresis).

RYANODEX® is associated with skeletal muscle weakness such as loss of grip strength and weakness in the legs, as well as drowsiness, dizziness, dysphagia, dyspnea, and decreased inspiratory capacity. Patients should not be permitted to ambulate without assistance until they have normal strength and balance. Care must be taken to prevent extravasation of RYANODEX® into the surrounding tissue due to the high pH of the reconstituted RYANODEX® suspension and potential for tissue necrosis.

Please see enclosed full Prescribing Information.

REFERENCES:

1. Riazi S, Larach MG, Hu C, Wijeyesundera D, Massey C, Kraeva N. Malignant hyperthermia in Canada: characteristics of index anesthetics in 129 malignant hyperthermia susceptible probands. *Anesth Analg*. 2014;118(2):381-387.
2. Larach MG, Gronert GA, Allen GC, Brandom BW, Lehman EB. Clinical presentation, treatment, and complications of malignant hyperthermia in North America from 1987 to 2006. *Anesth Analg*. 2010;110(2):498-507.
3. Riazi S, Kraeva N, Hopkins PM. Updated guide for the management of malignant hyperthermia. *Can J Anesth*. 2018;65(6):709-721.
4. Litman RS, Jones SB, Crowley M. Malignant hyperthermia: diagnosis and management of acute crisis Up to Date website. <https://www.uptodate.com/contents/malignant-hyperthermia-diagnosis-and-management-of-acute-crisis>. Accessed March 12, 2020.
5. Schütte JK, Becker S, Burmester S, et al. Comparison of the therapeutic effectiveness of a dantrolene sodium solution and a novel nanocrystalline suspension of dantrolene sodium in malignant hyperthermia susceptible pigs. *Eur J Anaesthesiol*. 2011;28(4):256-264.
6. RYANODEX® [prescribing information]. Woodcliff Lake, NJ: Eagle Pharmaceuticals, Inc.; 2017.
7. Data on file. Eagle Pharmaceuticals, Inc.
8. Dantrium Intravenous [prescribing information]. Chestnut Ridge, NY: Par Pharmaceutical, Inc.; 2016.
9. Revonto [prescribing information]. Louisville, KY: US WorldMeds, LLC; 2016.
10. Managing an MH crisis. Malignant Hyperthermia Association of the United States website. <http://www.mhaus.org/healthcare-professionals/managing-a-crisis>. Accessed March 12, 2020.



1 LIFE
at risk

1 VIAL
for most patients

1 PROVIDER
to reconstitute

1 MINUTE
to administer

1 NAME
to remember

EAGLE
PHARMACEUTICALS

© 2020 Eagle Pharmaceuticals, Inc. All rights reserved.
Woodcliff Lake, NJ 07677 MLR-RYM-US-0065

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use RYANODEX safely and effectively. See full prescribing information for RYANODEX.

RYANODEX® (dantrolene sodium) for injectable suspension, for intravenous use.

Initial U.S. Approval: 1974

INDICATIONS AND USAGE

RYANODEX is a skeletal muscle relaxant drug indicated for:

- Treatment of malignant hyperthermia in conjunction with appropriate supportive measures. (1)
- Prevention of malignant hyperthermia in patients at high risk. (1)

DOSAGE AND ADMINISTRATION

- **Treatment of Malignant Hyperthermia (MH)** (2.1)
 - Administer by intravenous push at a minimum of 1 mg/kg. If signs continue, administer additional intravenous boluses up to maximum cumulative dosage of 10 mg/kg.
 - Institute supportive measures (e.g., discontinue MH-triggering agents, manage metabolic acidosis, cooling if necessary, administer diuretics)
- **Prevention of MH in Patients at High Risk** (2.2)
 - Administer 2.5 mg/kg intravenously over a period of at least 1 minute, starting approximately 75 minutes prior to surgery.
 - Avoid agents that trigger MH.
 - Administer additional individualized doses during anesthesia and surgery if surgery is prolonged.
- **Pediatric Patients:** recommended weight-based dose is the same as for adults (2.3)
- **Reconstitution:** With 5 mL of sterile water for injection (without a bacteriostatic agent) prior to administration (2.4)

DOSAGE FORMS AND STRENGTHS

For injectable suspension: 250 mg of lyophilized powder in a single-dose vial for reconstitution (3)

CONTRAINDICATIONS

None (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage for Treatment of Malignant Hyperthermia
- 2.2 Dosage for Prevention of Malignant Hyperthermia
- 2.3 Dosage for Pediatric Patients
- 2.4 Reconstitution and Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Muscle Weakness
- 5.2 Somnolence and Dizziness
- 5.3 Potential for Tissue Necrosis with Extravasation

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Calcium Channel Blockers
- 7.2 Muscle Relaxants
- 7.3 Antipsychotics and Antianxiety Agents

WARNINGS AND PRECAUTIONS

- **Skeletal Muscle Weakness:** Ambulate patients with assistance until they have normal strength and balance (5.1)
- **Dyspnea, Respiratory Muscle Weakness, and Decreased Inspiratory Capacity:** Monitor patients for the adequacy of ventilation (5.1)
- **Dysphagia:** Assess patients for difficulty swallowing and choking (5.1)
- **Somnolence and Dizziness:** Can occur following RYANODEX administration; may persist up to 48 hours post-dose. Ambulate patients with assistance until they have normal strength and balance (5.2)
- **Tissue Necrosis with Extravasation:** Due to high pH of reconstituted RYANODEX for injectable suspension, care must be taken to prevent extravasation into surrounding tissues (5.3)

ADVERSE REACTIONS

- Administration to conscious subjects is associated with loss of grip strength and weakness in the legs, drowsiness and dizziness.
- Other common adverse reactions are: nausea, thrombophlebitis, tissue necrosis secondary to extravasation, urticaria and erythema, and injection site reactions (pain, erythema, swelling) (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Eagle at 1-855-318-2170 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Calcium Channel Blockers:** Concomitant use can cause cardiovascular collapse in association with marked hyperkalemia (7.1)
- **Muscle Relaxants:** Concomitant use with muscle relaxants may potentiate their effects (7.2)
- **Sedatives:** Concomitant use with sedative agents may potentiate their effects (7.3)

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** Based on animal data, may cause fetal harm (8.1)
- **Nursing Mothers:** Discontinue drug or nursing (8.3)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 09/2017

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

- 10.1 Overdosage Symptoms
- 10.2 Management of Overdosage

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

RYANODEX[®] is indicated for the:

- Treatment of malignant hyperthermia in conjunction with appropriate supportive measures [see *Dosage and Administration (2.1)*]
- Prevention of malignant hyperthermia in patients at high risk.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage for Treatment of Malignant Hyperthermia

In addition to RYANODEX treatment, institute the following supportive measures:

- Discontinue use of malignant hyperthermia (MH)-triggering anesthetic agents (i.e., volatile anesthetic gases and succinylcholine).
- Manage the metabolic acidosis
- Institute cooling when necessary
- Administer diuretics to prevent late kidney injury due to myoglobinuria (the amount of mannitol in RYANODEX is insufficient to maintain diuresis) [see *Description (11)*]

Administer RYANODEX by intravenous push at a minimum dose of 1 mg/kg. If the physiologic and metabolic abnormalities of MH continue, administer additional intravenous boluses up to the maximum cumulative dosage of 10 mg/kg.

If the physiologic and metabolic abnormalities reappear, repeat RYANODEX dosing by intravenous push starting with 1 mg/kg.

2.2 Dosage for Prevention of Malignant Hyperthermia

The recommended prophylactic dose of RYANODEX is 2.5 mg/kg administered intravenously over a period of at least 1 minute, starting approximately 75 minutes prior to surgery. Avoid agents that trigger MH.

If surgery is prolonged, administer additional individualized RYANODEX doses during anesthesia and surgery [see *Clinical Pharmacology (12.3)*].

2.3 Dosage for Pediatric Patients

The recommended weight-based dose of RYANODEX for pediatric patients in the treatment and prevention of MH is the same as for adults for these indications [see *Dosage and Administration (2.1, 2.2)*].

2.4 Reconstitution and Administration Instructions

The supplied lyophilized powder must be reconstituted prior to administration:

- (a) Reconstitute each vial of RYANODEX lyophilized powder by adding 5 mL of sterile water for injection (without a bacteriostatic agent). Do not reconstitute with any other solution (e.g., 5% dextrose injection, 0.9% sodium chloride injection).
- (b) Shake the vial to ensure an orange-colored uniform suspension. Visually inspect the vial for particulate matter and discoloration prior to administration.
- (c) Must use the contents of the vial within 6 hours after reconstitution. Store reconstituted suspensions at controlled room temperature (68°F to 77°F or 20°C to 25°C).

Do not dilute or transfer the reconstituted RYANODEX suspension to another container to infuse the product.

Administer the reconstituted RYANODEX suspension either:

- Into the intravenous catheter while an intravenous infusion of 0.9% sodium chloride injection, or 5% dextrose injection is freely running; or
- Into the indwelling catheter - after assuring its patency - without a freely running infusion. Flush the line to assure that there is no residual RYANODEX remaining in the catheter [*see Warnings and Precautions (5.3)*].

3 DOSAGE FORMS AND STRENGTHS

For injectable suspension: RYANODEX is a sterile, lyophilized powder containing 250 mg of dantrolene sodium for reconstitution, in single-dose vials

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Muscle Weakness

RYANODEX is associated with skeletal muscle weakness. The administration of RYANODEX in human volunteers has been associated with loss of grip strength and weakness in the legs. Patients should not be permitted to ambulate without assistance until they have normal strength and balance.

RYANODEX has been associated with dyspnea, respiratory muscle weakness, and decreased inspiratory capacity. Monitor patients for the adequacy of ventilation.

RYANODEX has been associated with dysphagia. Assess patients for difficulty swallowing and choking.

5.2 Somnolence and Dizziness

Somnolence and dizziness can occur following administration of RYANODEX and may persist up to 48 hours post-dose. Patients should not be permitted to ambulate without assistance until they have normal strength and balance. Patients must not operate an automobile or engage in other hazardous activities for 48 hours post-dose.

The concomitant use of sedative agents with RYANODEX may increase the risk of somnolence and dizziness.

5.3 Potential for Tissue Necrosis with Extravasation

Care must be taken to prevent extravasation of RYANODEX into the surrounding tissues due to the high pH of the reconstituted RYANODEX suspension and potential for tissue necrosis.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In a study designed to evaluate the safety and tolerability of RYANODEX, healthy volunteers were randomly assigned to receive treatment with RYANODEX or an active comparator at doses ranging from 1 mg/kg to 2.5 mg/kg.

- The RYANODEX dose was infused over the course of 1 minute for each of the doses evaluated.
- The active comparator was an injectable formulation of dantrolene sodium that differed from RYANODEX in that it contained dantrolene sodium and mannitol at concentrations of 0.33 mg/mL and 50 mg/mL, respectively, when reconstituted according to the product's prescribing information. The active comparator was infused at a rate that administered 20 mg of dantrolene per minute for each of the doses evaluated.

Table 1 displays the most common adverse events in this study. These data are not an adequate basis for comparison of the types or frequencies of adverse event types between RYANODEX and the dantrolene sodium comparator.

Adverse events increased in frequency with increasing doses in the trial, but did not differ in frequency between the two treatment groups. RYANODEX-treated subjects were more likely to report immediate adverse events of flushing, dystonia, and dysphagia than those receiving the active comparator.

In all dose groups, hand grip strength declined after dosing. In general, the decline in hand grip strength was more pronounced and occurred more rapidly in the RYANODEX-treated subjects in the 1.0, 1.75, 2.0 and 2.25 mg/kg treatment groups. In the 2.5 mg/kg treatment group, the decline in hand grip strength both in amount and duration was similar between the two treatment groups.

Table 1: Adverse Events in Healthy Volunteers

	Number of subjects (%)	
	RYANODEX [N=30] n (%)	Dantrolene Sodium Comparator [N=31] n (%)
Flushing	8 (27)	1 (3)
Somnolence	5 (17)	4 (13)
Dysphonia	4 (13)	1 (3)
Dysphagia	3 (10)	4 (13)
Nausea	3 (10)	3 (10)
Feeling abnormal	3 (10)	3 (10)
Headache	1 (3)	4 (13)
Vomiting	1 (3)	2 (6)
Vision blurred	1 (3)	1 (3)
Pain in extremity	1 (3)	1 (3)
Muscular weakness/Asthenia	1 (3)	1 (3)
Atrioventricular block	1 (3)	0
Tachycardia	1 (3)	0
Infusion site pain	1 (3)	0
Dizziness	1 (3)	0

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of another formulation of dantrolene sodium for injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Pulmonary Edema

There have been reports of pulmonary edema developing during the treatment of malignant hyperthermia crises with another dantrolene sodium dosage form. The contributory effect of the diluent volume and mannitol in these cases is not known.

Thrombophlebitis and Tissue Necrosis

There have been reports of thrombophlebitis following administration of intravenous dantrolene. Tissue necrosis secondary to extravasation has been reported [*see Warnings and Precautions (5.3)*].

Hypersensitivity/Anaphylactic Reactions

There have been reports of urticaria and erythema possibly associated with the administration of dantrolene sodium for injection. Anaphylaxis has been reported.

Injection Site Reactions

Injection site reactions including pain, erythema, and swelling, commonly due to extravasation, have been reported.

7 DRUG INTERACTIONS

7.1 Calcium Channel Blockers

Cardiovascular collapse in association with marked hyperkalemia has been reported in patients receiving dantrolene in combination with calcium channel blockers. The concomitant use of RYANODEX and calcium channel blockers is not recommended during the treatment of malignant hyperthermia.

7.2 Muscle Relaxants

The concomitant administration of RYANODEX with muscle relaxants may potentiate the neuromuscular block.

7.3 Antipsychotics and Antianxiety Agents

The concomitant administration of RYANODEX with antipsychotic and antianxiety agents may potentiate their effects on the central nervous system [*see Warnings and Precautions (5.2)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

Adequate and well controlled studies have not been conducted with RYANODEX in pregnant women. However, animal reproduction studies have been conducted with dantrolene sodium. In these studies, dantrolene sodium administered to rats and rabbits produced embryoletality (rabbits) and decreased pup survival (rats) at doses seven times the human oral dose.

RYANODEX should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.2 Labor and Delivery

In one uncontrolled study, 100 mg per day of prophylactic oral dantrolene sodium was administered to term pregnant patients awaiting labor and delivery. Dantrolene readily crossed the placenta, with maternal and fetal whole blood levels approximately equal at delivery; neonatal levels then fell approximately 50% per day for 2 days before declining sharply. No neonatal respiratory and neuromuscular side effects were observed in this study.

8.3 Nursing Mothers

Dantrolene is present in human milk. In one case report, low dantrolene concentrations (less than 2 micrograms per milliliter) were measured in the breast milk of a lactating woman during repeat intravenous dantrolene administration over 3 days. Because of the potential for serious adverse reactions of respiratory depression and muscle weakness in nursing infants from dantrolene, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

8.4 Pediatric Use

The safety and efficacy of RYANODEX in the treatment and prevention of malignant hyperthermia in pediatric patients is based on clinical experience with other intravenous dantrolene sodium products, which suggests adult weight-based doses are appropriate for pediatric patients.

8.5 Geriatric Use

Clinical studies of RYANODEX did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

10 OVERDOSAGE

10.1 Overdosage Symptoms

Overdosage symptoms include, but are not limited to, muscular weakness and alterations in the state of consciousness (e.g., lethargy, coma), vomiting, diarrhea, and crystalluria.

10.2 Management of Overdosage


Employ general supportive measures for acute overdosage of RYANODEX.

Administer intravenous fluids in fairly large quantities to avert the possibility of crystalluria. Maintain an adequate airway and keep artificial resuscitation equipment available. Institute electrocardiographic monitoring and carefully observe the patient. The value of dialysis in RYANODEX overdosage is not known.

11 DESCRIPTION

RYANODEX[®] (dantrolene sodium) for injectable suspension is a sterile lyophilized powder. RYANODEX is supplied in 20 mL vials containing 250 mg dantrolene sodium and the following inactive ingredients: 125 mg mannitol, 25 mg polysorbate 80, 4 mg povidone K12 and sufficient sodium hydroxide or hydrochloric acid for pH adjustment. When reconstituted with 5 mL sterile water for injection USP (without a bacteriostatic agent), this yields a suspension with a pH of approximately 10.3.

RYANODEX is a skeletal muscle relaxant. Chemically, RYANODEX is a hydrate of 1-[[[5-(4-nitrophenyl)-2-furanyl]methylene]amino]-2,4-imidazolidinedione sodium salt. The structural formula for the hydrated salt is:



The hydrated salt contains approximately 15% water (3-1/2 moles) and has a molecular weight of 399. The anhydrous salt (dantrolene) has a molecular weight of 336.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

In isolated nerve-muscle preparation, dantrolene has been shown to produce relaxation by affecting the contractile response of the muscle at a site beyond the myoneural junction. In skeletal muscle, dantrolene dissociates the excitation-contraction coupling, probably by interfering with the release of Ca^{++} from the sarcoplasmic reticulum.

In the anesthetic-induced malignant hyperthermia syndrome, evidence points to an intrinsic abnormality of skeletal muscle tissue. In affected humans, it has been postulated that “triggering agents” (e.g., general anesthetics and depolarizing neuromuscular blocking agents) produce a change within the cell which results in an elevated myoplasmic calcium. This elevated myoplasmic calcium activates acute cellular catabolic processes that cascade to the malignant hyperthermia crisis.

The addition of dantrolene to the “triggered” malignant hyperthermic muscle cell may reestablish a normal level of ionized calcium in the myoplasm. Inhibition of calcium release from the sarcoplasmic reticulum by dantrolene reestablishes the myoplasmic calcium equilibrium, increasing the percentage of bound calcium. In this way, physiologic, metabolic, and biochemical changes associated with the malignant hyperthermia crisis may be reversed or attenuated.

12.2 Pharmacodynamics

The administration of intravenous dantrolene to human volunteers was associated with loss of grip strength and weakness in the legs, as well as subjective CNS complaints [*see Warnings and Precautions (5.1)*].

12.3 Pharmacokinetics

The pharmacokinetics of RYANODEX was investigated in healthy volunteers following single-dose administration as an intravenous push over 60 seconds (dose range of 1 to 2.5 mg/kg), via a peripheral catheter. There was a dose-proportional increase in plasma exposure of dantrolene and its metabolite, 5-hydroxydantrolene. Table 2 presents pharmacokinetic parameters of dantrolene after administration of a single RYANODEX dose of 2.5 mg/kg. Time to peak dantrolene concentration was observed at the first time point collected (i.e., median T_{max} is 1 minute post-dose). The mean half-life ($t_{1/2}$) for dantrolene was independent of the RYANODEX dose administered and ranged from 8.5 hours to 11.4 hours over the 1 to 2.5 mg/kg dose range.

Table 2: Pharmacokinetic Parameters for Dantrolene in Healthy Volunteers (Mean \pm Standard Deviation)

RYANODEX dose ¹	C_{max} ($\mu\text{g}/\text{mL}$)	$\text{AUC}_{0-\text{inf}}$ ($\mu\text{g}\cdot\text{h}/\text{mL}$)	Half-life (hrs)	Clearance (L/hr)	Volume of distribution (L)
2.5 mg/kg	9.0 ± 4.6	77.7 ± 23.2	10.8 ± 2.2	2.5 ± 1.0	36.4 ± 11.7

¹ n=15; single-dose

When prophylactic intravenous dantrolene infusion was administered, whole blood dantrolene concentrations remained at a near steady state level for 3 or more hours after the infusion was completed.

Distribution

Based on assays of whole blood and plasma, slightly greater amounts of dantrolene are associated with red blood cells than with the plasma fraction of blood. Significant amounts of dantrolene are bound to plasma proteins, mostly albumin, and this binding is readily reversible. Binding to plasma protein is not significantly altered by diazepam, diphenylhydantoin, or phenylbutazone. Binding to plasma proteins is reduced by warfarin and clofibrate and increased by tolbutamide.

Information concerning the passage of dantrolene across the blood-brain barrier is not available.

Metabolism

Dantrolene is found in measurable amounts in blood and urine. Its major metabolites in body fluids are 5-hydroxy dantrolene and an acetylamino metabolite of dantrolene. Another metabolite with an unknown structure appears related to the latter. Dantrolene may also undergo hydrolysis and subsequent oxidation forming nitrophenylfuroic acid. Following RYANODEX 2.5 mg/kg dose administration in healthy volunteers, mean peak plasma concentration for the primary metabolite, 5-hydroxydantrolene, was 640 ng/mL, and was achieved by approximately 24 hours post-dose, with an average metabolite-to-parent exposure ratio of 0.27 ($AUC_{0-inf} = 21.2 \mu\text{g}\cdot\text{h/mL}$). Median $t_{1/2}$ for 5-hydroxydantrolene was 10 hours, the clearance was estimated to be 8.9 L/hr and the terminal phase volume of distribution was 144 L.

Dantrolene is metabolized by the liver, and it is possible that its metabolism may be enhanced by drugs known to induce hepatic microsomal enzymes. However, neither phenobarbital nor diazepam appears to affect dantrolene metabolism.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Sprague-Dawley female rats fed dantrolene sodium for 18 months at dosage levels of 15, 30, and 60 mg/kg/day showed an increased incidence of benign and malignant mammary tumors compared with concurrent controls. At the highest dose level (approximately the same as the maximum recommended daily human dose on a mg/m^2 basis), there was an increase in the incidence of benign hepatic lymphatic neoplasms. In a 30-month study in Sprague-Dawley rats fed dantrolene sodium, the highest dose level (approximately the same as the maximum recommended daily human dose on a mg/m^2 basis) produced a decrease in the time of onset of mammary neoplasms. Female rats at the highest dose level showed an increased incidence of hepatic lymphangiomas and hepatic angiosarcomas.

The only drug-related effect seen in a 30-month study in Fischer-344 rats was a dose-related reduction in the time of onset of mammary and testicular tumors. A 24-month study in HaM/ICR mice revealed no evidence of carcinogenic activity.

The significance of carcinogenicity data relative to use of RYANODEX in humans is unknown.

Mutagenesis

Dantrolene sodium has produced positive results in the Ames *S. Typhimurium* bacterial mutagenesis assay in the presence and absence of a liver activating system.

Impairment of Fertility

Dantrolene sodium administered to male and female rats at dose levels up to 45 mg/kg/day (approximately 1.4 times the maximum recommended daily human dose on a mg/m² basis) showed no adverse effects on fertility or general reproductive performance.

16 HOW SUPPLIED/STORAGE AND HANDLING

RYANODEX[®] (NDC 42367-540-32) is available in 20 mL vials containing a sterile lyophilized mixture of 250 mg dantrolene sodium for reconstitution with 5 mL sterile water for injection USP (without a bacteriostatic agent) to yield an orange colored injectable suspension.

Store unreconstituted product at 20 °C to 25 °C (68 °F to 77 °F) [see USP Controlled Room Temperature], with excursions permitted to 15 °C to 30 °C (59 °F to 86 °F) and avoid prolonged exposure to light.

17 PATIENT COUNSELING INFORMATION

Inform patients, their families, or their caregivers of the following:

Muscle Weakness

Muscle weakness (i.e., decrease in grip strength and weakness of leg muscles, especially walking down stairs) is likely to occur with the use of RYANODEX. Patients should be provided assistance with standing and walking until their strength has returned to normal [*see Warnings and Precautions (5.1)*].

Difficulty Swallowing

Caution is indicated at meals on the day of administration because difficulty swallowing and choking have occurred with the use of dantrolene sodium products in general; dysphagia has been reported with the use of RYANODEX [*see Warnings and Precautions (5.1)*].

Dizziness and Somnolence

The use of RYANODEX has been associated with dizziness and somnolence. [*see Warnings and Precautions (5.2)*].

Driving or Operating Machinery

Symptoms such as “lightheadedness” may occur. Since some of these symptoms may persist for up to 48 hours, patients must not operate an automobile or engage in other hazardous activity during this time [*see Warnings and Precautions (5.2)*].

Marketed by:

Eagle Pharmaceuticals, Inc.
Woodcliff Lake, NJ 07677